arXiv:1407.5789v1 [math.NT] 22 Jul 2014

A CURIOUS CONGRUENCE INVOLVING ALTERNATING HARMONIC SUMS

LIUQUAN WANG

Abstract. Let p be a prime and $\mathcal{P}_p$ the set of positive integers which are prime to p. We establish the following interesting congruence

$$\sum_{\substack{i+j+k=p^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk}\equiv\frac{p^{r-1}}{2}B_{p-3}\,(\mathrm{mod}\,p^r).$$

1. Introduction and Main Results

Let B_n be the n-th Bernoulli number, which is defined by

$$\frac{x}{e^x-1}=\sum_{n=0}^{\infty}\frac{B_n}{n}x^n.$$

In [4], Zhao found a curious congruence

$$\sum_{\substack{i+j+k=p\\ i,j,k>0}}\frac{1}{ijk}\equiv-2B_{p-3}\,(\mathrm{mod}\,p), \tag{1}$$

where $p\geq 3$ is a prime.

Let $\mathcal{P}_n$ denotes the set of positive integers which are prime to n. Recently, Wang and Cai [2] gave a generalization to this congruence, they have shown

$$\sum_{\substack{i+j+k=p^r\\ i,j,k\in\mathcal{P}_p}}\frac{1}{ijk}\equiv-2p^{r-1}B_{p-3}\,(\mathrm{mod}\,p^r).$$

For more variant and generalizations of (1), we refer the reader to see [3] and [5].

In this paper, we consider the alternating sums and obtain the following analogous result.

Theorem 1. *Let $p\geq 3$ be a prime and r a positive integer, then*

$$\sum_{\substack{i+j+k=p^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk}\equiv\frac{p^{r-1}}{2}B_{p-3}\,(\mathrm{mod}\,p^r). \tag{2}$$

Furthermore, we have

Date: June 2, 2014.

2010 *Mathematics Subject Classification.* Primary 11A07, 11A41.

Key words and phrases. congruences, Bernoulli numbers, harmonic sums.

Theorem 2. *Let n be a positive integer and $p \geq 3$ a prime, if $p^r | n$, $r \geq 1$ is an integer, then*

$$\sum_{\substack{i+j+k=n \\ i,j,k \in \mathcal{P}_p}} \frac{(-1)^i}{ijk} \equiv \frac{n}{2p} B_{p-3} \pmod{p^r}.$$

In particular, if $n = p^r$, Theorem 2 becomes Theorem 1.

2. Preliminaries

Lemma 1 (cf. Theorem 5.2 in [1])**.** *For any prime $p \geq 3$ we have*

$$\sum_{x=1}^{(p-1)/2} \frac{1}{x^3} \equiv -2B_{p-3} \pmod{p}.$$

The following Lemma is contained in [2], for the sake of completeness, we provide the proof here.

Lemma 2. *Let $p \geq 3$ be a prime and r a positive integer, for $1 \leq x \leq p-1$, define*

$$S(x, p^r) = \sum_{\substack{i \equiv x \pmod{p} \\ 1 \leq i \leq p^r - 1}} \frac{1}{i},$$

we have

$$S(x, p^r) \equiv p^{r-1} x^{-1} \pmod{p^r}.$$

Proof. For any integer $0 \leq k \leq p^r - 1$, write k as

$$k = a + p^{r-1} b, 0 \leq a \leq p^{r-1} - 1, 0 \leq b \leq p - 1.$$

Then

$$\begin{aligned} & S(x, p^{r+1}) - pS(x, p^r) \\ &= \sum_{k=0}^{p^r - 1} \frac{1}{x + kp} - p \sum_{k=0}^{p^{r-1} - 1} \frac{1}{x + kp} \\ &= \sum_{b=0}^{p-1} \sum_{a=0}^{p^{r-1} - 1} \frac{1}{x + (a + p^{r-1} b)p} - \sum_{b=0}^{p-1} \sum_{a=0}^{p^{r-1} - 1} \frac{1}{x + ap} \\ &= -p^r \sum_{b=1}^{p-1} \sum_{a=0}^{p^{r-1} - 1} \frac{b}{(x + ap)\{x + (a + p^{r-1} b)p\}}. \end{aligned} \tag{3}$$

Because

$$\sum_{b=0}^{p-1} b = \frac{p(p-1)}{2} \equiv 0 \pmod{p},$$

we have $S(x, p^{r+1}) - pS(x, p^r) \equiv 0 \pmod{p^{r+1}}$.

Note that $S(x, p) = x^{-1}$, by induction on r it's easy to see that $S(x, p^r) \equiv p^{r-1} x^{-1} \pmod{p^r}$. □

3. Proofs of the Theorems

Proof of Theorem 1. It's easy to see that

$$\begin{aligned}\sum_{\substack{i+j+k=p^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk}&=\sum_{\substack{i+j+k=p^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^{p^r-j-k}}{(p^r-j-k)jk}\\ &\equiv\sum_{\substack{j+k<p^r\\ j,k,j+k\in\mathcal{P}_p}}\frac{(-1)^{j+k}}{jk(j+k)}\\ &\equiv 2\sum_{\substack{1\le j<m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^m}{jm^2}\,(\mathrm{mod}\,p^r),\end{aligned}\tag{4}$$

where in the last equality we replace $j+k$ by m and used the fact

$$\frac{1}{jk(j+k)}=\frac{j+k}{jk(j+k)^2}=\big(\frac{1}{j}+\frac{1}{k}\big)\frac{1}{(j+k)^2}.$$

Note that

$$\sum_{\substack{1\le j<m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^m}{jm^2}=\sum_{\substack{1\le j<m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^{p^r-j}}{(p^r-m)(p^r-j)^2}\equiv\sum_{\substack{1\le j<m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^j}{mj^2}\,(\mathrm{mod}\ p^r).$$

Hence we have

$$\begin{aligned}2\sum_{\substack{1\le j<m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^m}{jm^2}&=\sum_{\substack{1\le j,m<p^r\\ j,m,m-j\in\mathcal{P}_p}}\frac{(-1)^m}{jm^2}\\ &=\sum_{\substack{1\le j,m<p^r\\ j,m\in\mathcal{P}_p}}\frac{(-1)^m}{jm^2}-\sum_{\substack{1\le j,m<p^r\\ j\in\mathcal{P}_p,j\equiv m\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{jm^2}\\ &=\Big(\sum_{\substack{1\le j<p^r\\ j\in\mathcal{P}_p}}\frac{1}{j}\Big)\Big(\sum_{\substack{1\le m<p^r\\ m\in\mathcal{P}_p}}\frac{(-1)^m}{m^2}\Big)-\sum_{\substack{1\le j,m<p^r\\ j\in\mathcal{P}_p,j\equiv m\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{jm^2}\\ &\equiv-\sum_{\substack{1\le j,m<p^r\\ j\in\mathcal{P}_p,j\equiv m\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{jm^2}\,(\mathrm{mod}\,p^r),\end{aligned}\tag{5}$$

here the last congruence equality follows from the fact

$$\sum_{\substack{1\le j<p^r\\ j\in\mathcal{P}_p}}\frac{1}{j}=\frac{1}{2}\sum_{\substack{1\le j<p^r\\ j\in\mathcal{P}_p}}\big(\frac{1}{j}+\frac{1}{p^r-j}\big)=\frac{1}{2}\sum_{\substack{1\le j<p^r\\ j\in\mathcal{P}_p}}\frac{p^r}{j(p^r-j)}\equiv 0\,(\mathrm{mod}\,p^r).$$

By Lemma 1 we deduce that

$$\sum_{x=1}^{p-1}\frac{(-1)^x}{x^3}=-\sum_{x=1}^{p-1}\frac{1}{x^3}+2\sum_{x=1}^{\frac{p-1}{2}}\frac{1}{(2x)^3}\equiv-\frac{1}{2}B_{p-3}\,(\mathrm{mod}\,p).$$

Thus by Lemma 2 we have

$$\begin{aligned}\sum_{\substack{1\le j,m<p^r\\ j\in\mathcal{P}_p, j\equiv m\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{jm^2}&=\sum_{x=1}^{p-1}\Big(\sum_{\substack{1\le j<p^r\\ j\equiv x\,(\mathrm{mod}\ p)}}\frac{1}{j}\Big)\Big(\sum_{\substack{1\le m<p^r\\ m\equiv x\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{m^2}\Big)\\ &=p^{r-1}\sum_{x=1}^{p-1}\frac{1}{x}\sum_{\substack{1\le m<p^r\\ m\equiv x\,(\mathrm{mod}\ p)}}\frac{(-1)^m}{m^2}\\ &=p^{r-1}\sum_{x=1}^{p-1}\frac{1}{x}\sum_{a=0}^{p^{r-1}-1}\frac{(-1)^{x+ap}}{(x+ap)^2}\\ &\equiv p^{r-1}\sum_{x=1}^{p-1}\frac{(-1)^x}{x^3}\\ &\equiv -\frac{p^{r-1}}{2}B_{p-3}\,(\mathrm{mod}\,p^r).\end{aligned}\tag{6}$$

Combining (4), (5) and (6), we complete the proof of Theorem 1. □

Proof of Theorem 2. Let $n=p^rm$, we only need to show

$$\sum_{\substack{i+j+k=mp^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk}\equiv\frac{mp^{r-1}}{2}B_{p-3}\,(\mathrm{mod}\,p^r).\tag{7}$$

For every triple (i,j,k) of positive integers which satisfies $i+j+k=mp^r, i,j,k\in\mathcal{P}_p$, we rewrite

$$i=xp^r+i_0, j=yp^r+j_0, k=zp^r+k_0,$$

where $1\le i_0,j_0,k_0\le p^r-1$ are prime to p and x,y,z are nonnegative integers.

Since $3\le i_0+j_0+k_0<3p^r$ and $i_0+j_0+k_0=(m-x-y-z)p^r$, it's easy to see that

$$\begin{cases} i_0+j_0+k_0=p^r\\ x+y+z=m-1\end{cases}\quad\text{or}\quad\begin{cases} i_0+j_0+k_0=2p^r\\ x+y+z=m-2\end{cases}.$$

Note that

$$\sum_{\substack{x+y+z=m-1\\ x,y,z\ge0}}(-1)^x=\sum_{k=0}^{m-1}(-1)^k(m-k)=\big[\frac{m+1}{2}\big],$$

$$\sum_{\substack{x+y+z=m-2\\ x,y,z\ge0}}(-1)^x=\sum_{k=0}^{m-2}(-1)^k(m-1-k)=\big[\frac{m}{2}\big],$$

here $[x]$ denotes the integer part of x. Hence

(8)
$$\begin{aligned}\sum_{\substack{i+j+k=mp^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk} &= \sum_{\substack{i_0+j_0+k_0=p^r\\ x+y+z=m-1\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{(-1)^{xp^r+i_0}}{(xp^r+i_0)(yp^r+j_0)(zp^r+k_0)}\\ &\quad+\sum_{\substack{i_0'+j_0'+k_0'=2p^r\\ x+y+z=m-2\\ i_0',j_0',k_0'\in\mathcal{P}_p\\ i_0',j_0',k_0'<p^r}}\frac{(-1)^{xp^r+i_0'}}{(xp^r+i_0')(yp^r+j_0')(zp^r+k_0')}\\ &\equiv \sum_{\substack{i_0+j_0+k_0=p^r\\ x+y+z=m-1\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{(-1)^{x+i_0}}{i_0j_0k_0}+\sum_{\substack{i_0'+j_0'+k_0'=2p^r\\ x+y+z=m-2\\ i_0',j_0',k_0'\in\mathcal{P}_p\\ i_0',j_0',k_0'<p^r}}\frac{(-1)^{x+i_0'}}{i_0'j_0'k_0'}\\ &\equiv \Big[\frac{m+1}{2}\Big]\sum_{\substack{i_0+j_0+k_0=p^r\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{1}{i_0j_0k_0}+\Big[\frac{m}{2}\Big]\sum_{\substack{i_0'+j_0'+k_0'=2p^r\\ i_0',j_0',k_0'\in\mathcal{P}_p\\ i_0',j_0',k_0'<p^r}}\frac{(-1)^{i_0'}}{i_0'j_0'k_0'}\pmod{p^r}.\end{aligned}$$

For the second sum in (8), since $(i_0,j_0,k_0)\leftrightarrow(p^r-i_0,p^r-j_0,p^r-k_0)$ gives a bijection between the solutions of $i_0+j_0+k_0=p^r$ and $i_0'+j_0'+k_0'=2p^r$ (here $i_0',j_0',k_0'<p^r$), we have

(9)
$$\begin{aligned}\sum_{\substack{i_0'+j_0'+k_0'=2p^r\\ i_0',j_0',k_0'\in\mathcal{P}_p\\ i_0',j_0',k_0'<p^r}}\frac{(-1)^{i_0'}}{i_0'j_0'k_0'} &= \sum_{\substack{i_0+j_0+k_0=p^r\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{(-1)^{p^r-i_0}}{(p^r-i_0)(p^r-j_0)(p^r-k_0)}\\ &\equiv \sum_{\substack{i_0+j_0+k_0=p^r\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{(-1)^{i_0}}{i_0j_0k_0}\pmod{p^r}.\end{aligned}$$

Combining (8) with (9) and apply Theorem 1, we deduce

$$\begin{aligned}\sum_{\substack{i+j+k=mp^r\\ i,j,k\in\mathcal{P}_p}}\frac{(-1)^i}{ijk} &= \Big(\Big[\frac{m+1}{2}\Big]+\Big[\frac{m}{2}\Big]\Big)\sum_{\substack{i_0+j_0+k_0=p^r\\ i_0,j_0,k_0\in\mathcal{P}_p}}\frac{(-1)^{i_0}}{i_0j_0k_0}\\ &\equiv \frac{mp^{r-1}}{2}B_{p-3}\pmod{p^r},\end{aligned}$$

hence (7) is true. □

4. Concluding Remarks

We leave two open questions for future research.

Question 1. Can we find an arithmetical function $f(n)$ such that

$$\sum_{\substack{i+j+k=n\\ i,j,k\in\mathcal{P}_n}}\frac{(-1)^i}{ijk}\equiv f(n)\pmod{n}.$$

Question 2. For $n \geq 4$, can we find some analogous result for the following sum

$$\sum_{\substack{i_1+i_2+\cdots+i_n=p^r \\ i_1,i_2,\cdots,i_n\in\mathcal{P}_p}} \frac{(-1)^{i_1}}{i_1 i_2\cdots i_n}$$

modulo p^r?

References

[1] Z. Sun, Congruence concerning Bernoulli numbers and Bernoulli polynomials, Discrete Appl. Math, 105 (2000), 193-223.
[2] L. Wang and T. Cai, A curious congruence modulo prime powers, J. Number Theory, 144 (2014), 15-24.
[3] B. Xia and T. Cai, Bernoulli numbers and congurence for harmonic sums, Int. J. Number Theory, 06 (2010), 849-855.
[4] J. Zhao, Bernoulli numbers, Wolstenholme's theorem, and p^5 variations of Lucas' theorem, J. Number Theory, 123 (2007), 18-26.
[5] X. Zhou and T. Cai, A generalization of a curious congruence on harmonic sums, Proc. Amer. Math. Soc., 135 (2007), 1329-1333.

Department of Mathematics, National University of Singapore, Singapore, 119076, Singapore
E-mail address: mathlqwang@163.com